

MENU



Home > Books > Complementary Therapies



## CHAPTER METRICS OVERVIEW

97 Chapter Downloads

[View Full Metrics](#)



OPEN ACCESS PEER-REVIEWED CHAPTER

# An Introduction to Chiropractic BioPhysics® (CBP®) Technique: A Full Spine Rehabilitation Approach to Reducing Spine Deformities

Submitted: October 13th, 2021, Reviewed: January 16th, 2022, Published: May 19th, 2022  
IntechOpen uses cookies to offer you the best online experience. By continuing to use our site, you agree to

DOI: 10.5772/intechopen.102686

WRITTEN BY

Got it!

Deed E. Harrison and Paul A. Oakley



REGISTER TO DOWNLOAD FOR FREE



Share



Cite



FROM THE EDITED VOLUME

# Complementary Therapies

Edited by Mario Bernardo-Filho,  
Redha Tair, Danúbia da Cunha de  
Sá-Caputo and Adérito Seixas

[Book Details](#)

[Order Print](#)

ADVERTISEMENT

## Abstract

Chiropractic Biophysics® (CBP®) technique is a full-spine and posture correcting method that incorporates mathematical principles into a unique approach to treat spinal disorders. It considers that the identification of postural rotations and translations of human postures are first evaluated and compared to the radiographic assessment of the spine alignment. Mirror image® postural positions and movements are utilized including spinal extension positions to improve the spine and posture towards a normal/ideal alignment. Specifically, corrective exercises, corrective traction and chiropractic adjustments are performed encompassing a multimodal rehabilitation program with the goal of improving the posture and spine alignment. CBP Rehabilitation programs are typically performed in-office with supportive at-home measures. Repeat assessment including radiographs are used to quantify and monitor structural improvements. CBP technique is an evidence-based approach to treat spine deformities and is supported by all forms of clinical evidence including systematic literature reviews, randomized controlled trials, non-randomized controlled trials, case reports/series as well as in

trials, case reports/series as well as is supported by biomechanical posture-spine coupling validity, radiographic and posture analysis reliability/repeatability and use of a validated biomechanical spinal model as the outcome goal of care. CBP technique is a proven method to improve pain, disability and quality of life in those with structural deformities.

## Keywords

spine deformity

structural rehabilitation

traction

exercise

chiropractic

Author Information

Show +

## Chapter sections

# 1. Introduction

Chiropractic Biophysics<sup>®</sup> (CBP<sup>®</sup>) technique is a full-spine and posture correcting method that incorporates engineering and mathematical principles into a unique

approach in the treatment of spine disorders [1, 2, 3, 4, 5]. CBP technique is best described as a 'structural' rehabilitation approach as opposed to 'functional' rehabilitation that typically encompasses physiotherapeutic modalities, stretching and exercises to regain function. The goal in structural rehabilitation is to restore the spine alignment and posture to as near normal as possible.

CBP operates on three main premises: 1. There is a normal/ideal static spinal configuration; 2. Abnormal alterations of the spine/posture result in abnormal function disrupting homeostatic balance; 3. Altered static spine/postural alignment results in abnormal dynamics [1]. The contemporary spine literature supports all three of these premises (See Section 4). CBP technique has published research on many facets of the technique including defining what normal/ideal spine alignment is, how to measure spine alignment parameters with reliable and repeatable methods, how to correct/re-align spinal displacements, and evidence proving correcting spine and postural displacements correlates with improvements in pain, disability and quality

improvements in pain, disability and quality of life (QOL) measures (These studies are detailed later).

Herein, an overview is given of the scientific approach to treating spine disorders (i.e. subluxation) by the unique approach of CBP technique. A review will be given of the historical beginnings of CBP, rotations and translations of posture, the Harrison normal spinal model, radiographic analysis, posture and spinal coupling, the CBP protocol, clinical evidence of efficacy as well as the safety of the use of X-rays (The term 'X-rays' imply the use of plain radiographs throughout this chapter).

ADVERTISEMENT

No Evidence Found That E-Ciga...




## 2. Historical beginnings

Donald D. Harrison, who had a Master's degree in Mechanical Engineering and a Doctorate degree in Applied Mathematics developed a devote urgency to bring contemporary science to chiropractic. In the late 1970s, Harrison was the main instructor for the chiropractic technique named 'Pettibon.' Dissatisfied with the failure to produce spinal correction, he often incorporated his own methods in certain cases to better attain spine and posture improvements. It was in the treatment of one particular case (circa 1980) where he discovered that the body must be treated using the principles of mathematics; the term 'mirror image<sup>®</sup>' adjusting he later coined to describe these new approaches [1].

A 1974 paper by Panjabi et al. describes a Cartesian coordinate system for use in the description and study of joint biomechanics (Figure 1) [6]. Harrison was the first to apply this system of analysis to upright human posture (Figures 2 and 3). Harrison began

discovering the rotations and translations of human posture in 1980. During the early 1980s, the analysis system evolved to incorporate a full spine analysis of the head, rib cage and pelvis in three-dimensions. The technique methods continued to evolve with intellectual contributions from early practitioners of CBP including among others, Drs. DeGeorge, Gambale, Pope and Deed Harrison (founder's son).




**Figure 1.**

A vertebra described in terms of rotations about and translations along the x, y, and z-axes on a cartesian coordinate system as proposed by





Cartesian coordinate system as proposed by Panjabi (courtesy CBP seminars).



**Figure 2.**

If the head, thoracic cage, and pelvis are considered rigid bodies, then the possible rotations in three-dimensions are illustrated. Flexion and extension are rotations on the x-axis, axial rotation is about the y-axis, and lateral flexion is rotation about the z-axis (courtesy CBP seminars).






### Figure 3.

If the head, thoracic cage, and pelvis are considered rigid bodies, then the possible translations in three-dimensions are illustrated. Lateral translations occur along the x-axis, vertical translations occur along the y-axis, and anterior-posterior translations (protraction-retraction) occurs along the z-axis (courtesy CBP seminars).

One of the unique methods within CBP is the use of 'extension traction' to restore the normal cervical or lumbar lordosis (Figures 4 and 5). The first cervical extension traction was with use of an inclined bench that utilized a camlock and pulley system to hyperextend the neck by pulling on the


hyperextend the neck by pulling on the forehead [7]. This is the traction used in the first CBP non-randomized controlled clinical trial (nRCT) that showed that no traction either by no treatment or only cervical manipulation but no traction resulted in no improved alignment, while the traction group (also receiving cervical spinal manipulation) achieved improved lordosis [7].



**Figure 4.**

Cervical extension traction (courtesy CBP seminars).





**Figure 5.**

Lumbar extension traction (courtesy CBP seminars).

Further development in cervical traction involved the addition of a posterior-to-anterior (PA) pull through the mid cervical spine with simultaneous extension and distraction of the head while sitting in a chair, so-called ‘Pope’s 2-way’ traction (Figure 4) [8]. A slight modification of this traction involves the use of a chin-forehead strap to add weight directly to the patients head as an extension-compression 2-way traction (Figure 4) [9]. More recently, a cervical extension orthotic (Denneroll) has been shown to be effective at increasing cervical lordosis (Figure 4).

In the mid 1990s, Deed Harrison helped to

develop precision vectors for lumbar extension traction (Figure 5), where the first nRCT showing lumbar curve restoration was published in the *Archives of Physical Medicine and Rehabilitation* in 2002 and concluded: “This new method of lumbar extension traction is the first nonsurgical rehabilitative procedure to show increases in lumbar lordosis in chronic LBP (*low back pain*) subjects with hypolordosis” [10]. A lumbar extension orthotic device by Denneroll is also used for lumbar extension traction (Figure 5).

CBP technique is one of the most scientifically based posture and spine correcting techniques. There are many randomized controlled trials (RCTs), nRCTs, and well over 100 case reports/series documenting the improvement of diverse spine deformity patterns with concomitant reduction of pain, disability and increased QOL measures [11].

---

### 3. Rotations and translations of posture

The main strength of CBP technique is its fundamental underpinnings in engineering and mathematics [1]. It is a general theorem that any object can be decomposed as a rotation, a translation and a deformation [12]. Acknowledging that deformation of living tissues occurs, as in compressing of discs, ligaments, muscles etc., we divert attention to rotations and translations of posture. The main masses of the body, namely the head, thorax and pelvis can be described in relation to the body mass below within a Cartesian coordinate system (Figures 2 and 3). That is, the head is described in relation to the thorax, the thorax in relation to the pelvis, and the pelvis in relation to the feet [1, 13].

Any rotations or translations of the body masses as seen in neutral posture via external observation or internally by X-ray is acknowledged as abnormal. Therefore, no

offset of the masses equates to the normal postural alignment (i.e. un-subluxated position). It is important to note that in the assessment of a patient, it is the presence of a rotation or translation in the neutral standing position that is abnormal. When Harrison first applied this method of analysis, the treatment became apparent with the postural diagnosis. That is, for any rotation or translation apparent in neutral standing posture, the opposite position would need to be the treatment as applied during exercises, spinal traction or spinal adjustments, as this is the mathematical solution, “the exact reversing of the patient’s abnormal posture” [1]. In fact, because the soft tissues require a significant magnitude of stress and strains to attempt to correct the spinal position via mirror image methods, Harrison suggested that postural reflections (i.e. ‘mirror image’ adjustments) need to be applied in “twice the negative of the translation distances and rotation angles” [1].


It should be noted when Harrison finally developed the full spine analysis of rotations and translations of posture in the mid 1980s, he discovered that virtually 50% of all



he discovered that virtually 50% of all human movements had never been studied (except forward head posture). Thus, the Harrison research group performed several studies to evaluate the normal range of motion for several translation postures including lateral head and thoracic postures as well as anterior and posterior thoracic translation postures (Discussed in Section 6) [2, 3]. Clinically, the spinal coupling patterns as discovered to be associated with these common postural positions are of utmost importance in the treatment of these spinal disorders.

Importance of the study of these never previously studied translation postures can be highlighted in the distinction between true scoliosis and 'pseudo-scoliosis' (Figure 6) [14]. Pseudo-scoliosis is a lateral thoracic translation posture that characteristically features little to no vertebral rotation (simple to correct) [15, 16], whereas, true scoliosis characteristically features significant vertebral rotation (and is typically much more difficult to treat). X-ray screening of the spine is the only way to differentiate true scoliosis from pseudo-scoliosis.






## Figure 6.


Posture image and antero-posterior lumbar radiographs depicting a left lateral thoracic translation (side shift). Both patients in the radiographs have a 20 mm left lateral shift of T10 off midline. Left patient has a pure left lateral thoracic translation posture, aka ‘pseudo-scoliosis.’ Right patient has a true left lumbar scoliosis (vertebral rotation). Green line is vertical; red line highlights patient alignment (courtesy CBP seminars).

As mentioned, the absence of rotations and translations of the body masses in standing posture is normal. However, the shape of the spine position, particularly in the sagittal plane has traditionally been debated.

## 4. The Harrison normal spine model

In the mid 1990s to the mid 2000s, the Harrison research team performed a series of spine modeling studies of the sagittal spinal curves (Figure 7) [17, 18, 19, 20, 21, 22, 23, 24]. To this day, this seminal work serves as the treatment outcome goal (i.e. gold standard) for providing structural rehabilitation by CBP methods (Figure 8). In a series of systematic studies, elliptical shape modeling of the path of the posterior longitudinal ligament was performed as it could be easily compared to the posterior vertebral body margins on X-rays, the same anatomical region used for measuring the sagittal spinal curves (i.e. Harrison posterior tangents (Figure 9) [25, 26, 27, 28]).






**Figure 7.**

The Harrison normal sagittal spine model as the path of the posterior longitudinal ligament. The cervical, thoracic and lumbar curves are all portions of an elliptical curve having a unique minor-to-major axis ratio. The cervical curve is circular meaning the minor and major axes are


equal (courtesy CBP seminars).



## Figure 8.

Three patients demonstrating dramatically different spine alignment patterns. Left: excessive lumbar hyperlordosis, L4 anterolisthesis, and excessive anterior sagittal balance in a mid-aged female with disabling low back pain; middle: excessive thoracolumbar kyphosis and early degenerative changes in a mid-aged male; right: excessive thoracic hyperkyphosis in a young male with Scheuermann's disease. Red line is contiguous with posterior vertebral body margins; green line represents Harrison normal spinal model

represents Harrison normal spinal model  
(courtesy PAO).



**Figure 9.**

Harrison posterior tangent method involves lines drawn contiguous with the posterior vertebral body margins. Intersegmental as well as regional sagittal curves are easily quantified having a standard way of measurement with in about 2°

standard error of measurement within about 2° (courtesy CBP seminars).

Computer iterations of spine shape modeling were applied to determine the best-fit geometric spinal shapes by fitting ellipses of varying minor-to-major axis ratios to the digitized data points from the posterior vertebral body corners from X-ray samples for each of the three regions of the spine (cervical [[17](#), [18](#), [19](#)], thoracic [[20](#), [21](#)], and lumbar spine [[22](#), [23](#), [24](#)]). As shown in [Figure 7](#), the Harrison normal spinal model features a circular cervical lordosis, an elliptical thoracic curve featuring greater curvature cephalad with a straightened thoraco-lumbar junction and an elliptical lumbar lordosis showing a greater distal lumbar curvature. The spine is assumed to be vertical in the front view.

Although some have attempted to criticize the Harrison normal spinal model, it is important to acknowledge that it has been validated in several ways. Simple analysis of alignment data on samples of normal, asymptomatic populations have been done [[17](#), [18](#), [19](#), [20](#), [21](#), [22](#), [23](#), [24](#)]. Comparison studies between normal samples to

symptomatic samples have been performed [17, 29]. Comparisons between normal samples to theoretical ideal models have been done [17, 18, 20, 23]. Statistical differentiation of asymptomatic subjects from symptomatic pain group patients based on alignment data has been performed [19, 24].

In subsequent biomechanical modeling studies, the Harrison group used a validated postural loading model to verify that sagittal spinal balance and the sagittal curves of the spine are critical biomechanical parameters for maintaining postural load balance in healthy subjects [30]. Keller et al. [30] stated “because the pattern of [intervertebral disc] IVD postural stresses mirrored the sagittal curvatures and sagittal displacement of the spine, a failure of the IVD’s hydrostatic mechanism under these sustained loads could occur”. In a similar biomechanical modeling study, Harrison et al. determined that anterior sagittal thoracic posture (anterior thorax translation relative to the pelvis) resulted in significant increases in disc loads and stresses for all vertebral levels below T9 and that the extensor muscle loads required to maintain static equilibrium in



required to maintain static equilibrium in upright anterior posture increased almost five times that of normal [31]. In another study Keller et al. [32] determined that “postural forces are responsible for initiation of osteoporotic spinal deformity in elderly subjects”.

The Harrison group also used an elliptical shell model to evaluate the loads and bending moments on the cervical vertebrae in varying cervical spine deformity alignments [33, 34]. They found that in normal lordosis the anterior and posterior vertebral body stresses are nearly uniform and minimal, whereas, in cervical deformity configurations having kyphosis (S-shape kyphosis high or low, total kyphosis), the vertebral body stresses are ‘very large’ and opposite in direction compared to normal lordosis [33]. They concluded “This analysis provides the basis for the formation of osteophytes (Wolff’s Law) on the anterior margins of vertebrae in kyphotic regions of the sagittal cervical curve. This indicates that any kyphosis is an undesirable configuration in the cervical spine” [33]. Anterior head translation and a ‘military’ neck also displayed significantly increased vertebral



body stresses that are reverse in direction from C5-T1 and are also proven to be “undesirable configurations in the cervical spine” [34].

ADVERTISEMENT

---

## 5. Radiographic analysis

All radiographs should be taken in the ‘neutral’ standing position with the feet positioned with the heels at hips width apart. This is to avoid any induced postural deviations due to foot position. Also, to ensure a reproducible neutral (i.e. natural) body position, the subject should close their eyes and nod the head back and forth a couple times to where the subject should stop in their preferred position and then open their eyes while maintaining this adopted stance. Any postural misalignments seen in the subject should not be corrected. The lower body mass on the particular view being

lower body mass on the particular view being taken should be centered to the bucky. All X-rays should be taken without footwear.


It should be mentioned that the measurement of different sagittal spinal contours including regional curves or absolute rotation angles (ARAs) (i.e. cervical/lumbar lordosis; thoracic kyphosis) and intersegmental relative rotation angles (RRAs) between adjacent vertebrae can be easily quantified by use of the Harrison posterior tangent (HPT) lines (Figure 9) [25, 26, 27, 28]. The HPT method is preferred for three main reasons, 1. The posterior margins of the vertebral bodies are less affected by osteoarthritic changes as compared to the anterior margins which makes anatomical measurements more reliable and valid; 2. The posterior tangents are contiguous with the slope of the spinal curves and represent the first derivative in an engineering analysis and therefore, their intersection accurately depicts the sagittal configuration; 3. The HPT method has a small standard error of measurement (SEM) of approximately  $2^\circ$  versus higher SEMs with the Cobb ( $4.5\text{--}10^\circ$ ) [25, 26, 27]. This is why the HPT method is superior to other methods of sagittal spine

measurement including the popular Cobb method.

Generally, the global curves are measured as C2-C7, T1-T12, and L1-L5, however since the inflection of the cervical lordosis to thoracic kyphosis occurs at T1, some clinicians prefer to measure the cervical curve from C1-T1, and the thoracic curve from T2-T11 or T3-T10. Anterior sagittal translation distances are simply measured by the horizontal displacement offset between comparison vertebrae such as C2-S1, C2-C7 or T1, T1-T12, etc.

The anterior-to-posterior (AP) or PA X-rays are taken using the same postural positioning. The modified Risser-Ferguson method is employed to measure coronal plane alignment (Figure 10) [28]. On the AP/PA cervicothoracic view an upper angle is created as the angle between the best fit line of the upper cervical segments and intersection with the bite line, and a lower angle is formed between the best fit lines of the upper to lower spine segments [28]. The Rz angle is the angle formed by a vertical axis line (VAL) drawn from T4 and the lower cervicothoracic best fit line. Normal upper

cervicothoracic best fit line. Normal upper angle, lower angle and Rz cervicothoracic angles are  $90^\circ$ ,  $0^\circ$  and  $0^\circ$ , respectively. The AP/PA thoracic view may show an angle. The lumbo-pelvic view has an upper angle, the angle between the best fit line of the upper versus lower lumbar segments, and a lower angle, the angle between the best fit line between the lower segments and the horizontal pelvic line [28]. The upper angle and lower angle should be  $0^\circ$  and  $90^\circ$ , respectively. Any regional or full-spine coronal balance offset (i.e. imbalance) can be easily quantified as the horizontal distance between the uppermost segment to the lowermost segment (e.g. C2-T2, T1-T12, T12-S1, C2-S1).



**Figure 10.**

AP radiographic line drawing by modified Risser-Ferguson method.

ADVERTISEMENT

---

## 6. Posture and spinal coupling


Postural rotations and translations as described by Harrison (Figures 2 and 3) are understood as ‘main motions’ and the corresponding spinal displacements to accommodate the postural positions are termed ‘coupled motions’ [2, 3, 35, 36, 37, 38]. In CBP, a considerable clinical significance is placed on the correlation between the patient’s three-dimensional postural presentation (posture displacement in terms of rotations and translations) and the two-dimensional X-ray coupled motion (spinal rotations and translations) [2, 3, 38].

Of prime importance is the appreciation that unless there is buckling, anomalies or ligament damage, standing neutral postural

rotation and translation displacements of the head or thorax *cause* the vertebral spinal coupling patterns as seen on X-ray. If a patient's rotations and/or translations of posture 'match' the associated spinal coupling pattern as expected (i.e. normal coupling), then it is considered an 'easy' or typical case and the intuitive mirror image application of CBP methods would apply. When the patient's rotations and/or translations of posture do not match the expected spine coupling pattern (i.e. spinal coupling does not match postural displacement), then it is considered an atypical case where the clinician needs to consider alternative (i.e. more complicated) strategies for spine rehabilitation.

A classic demonstration of the 'matching' versus 'mismatching' of rotations and translations of posture and spine coupling patterns can be illustrated with forward head posture, aka, anterior head translation (AHT) (Figure 11). The natural and expected spine coupling with a forward translated head posture involves lower cervical spine flexion and upper cervical spine extension. As seen in Figure 11, many spine different vertebral coupling patterns are possible

vertebral coupling patterns are possible including hyperlordosis, hypolordosis, or kyphosis and accordingly, each cervical configuration requires its own unique application of CBP methods for its ideal correction.



**Figure 11.**

Forward head translation as shown in posture and in three unique lateral cervical radiographs. All three X-ray images have about 25 mm of forward head translation. Left: hyperlordosis; middle: hypolordosis; right: kyphosis. Green line is normal alignment; red line highlights patient alignment.

These cervical spine patterns have been termed harmonics and their presence can only be determined by radiography [2, 39]. Importantly, in CBP treatment approaches, each cervical spine coupling pattern (harmonic) requires its own unique treatment protocol. This is why many

manual therapy approaches (e.g. Mackenzie head retractions) are inadequate at correcting posture and spine alignment as these are prescribed universally (i.e. 'blackbox treatment') resulting in many patients receiving treatment protocols that are contraindicated. A patient with a hyperlordotic cervical spine should never be prescribed neck extension exercises as this would dynamically hyperextend the cervical joints. A patient with a complete cervical kyphosis should never be prescribed head retraction exercises as this often 'buckles' the spine into further kyphosis.

Also, as mentioned and illustrated in Figure 6, 'pseudo-scoliosis' or pure lateral translations of the thorax (or head) must be distinguished from true scoliosis by examination of the spinal coupling patterns [14]. If there is minimal or no vertebral rotation then this represents a typical case requiring CBP mirror image postural correction [3]. If there is vertebral rotation then it is considered true scoliosis and a completely different application of CBP methods (i.e. non-commutative properties of finite rotation angles [40, 41]). Case examples of the special application of CBP



methods in the treatment of scoliosis is described later.


ADVERTISEMENT

---

## 7. CBP protocol

The CBP patient management protocol [2, 3, 4] involves all typical initial patient examination procedures including the consultation, examination as well as pain, disability and quality of life questionnaires (Figure 12). In addition, CBP treatment consideration requires, without exception, a full-spine posture assessment as well as full-spine AP and lateral standing radiographs. Posture needs to be either qualitatively, but ideally quantitatively assessed as rotations and translations of the head, thorax and pelvis in three-dimensions (Figures 2 and 3). The X-rays need to be digitized and quantified, ideally with the Harrison posterior tangent method for the sagittal images and with the modified Risser-

# Ferguson on the AP images.



**Figure 12.**

CBP protocol treatment algorithm.

As seen in Figure 12, if appropriate, a new patient should be treated for their acute pain that is distinct and separate from CBP

that is distinct and separate from CBT methods. It is recommended that the acute 'pain care' treatment include spinal manipulation, stretching (e.g. proprioceptive neuromuscular facilitation (PNF), Yoga, etc.), heat/ice, soft tissue myofascial therapy (e.g. transverse friction, Nimmo-receptor tonus technique, etc.). Once the patient experiences some initial pain relief (e.g. 6–12 treatments) they can be re-assessed and graduated to CBP structural rehabilitation. The decision to first treat a new patient with 'acute' pain care is a clinical decision that is mainly for patients that have either never seen a chiropractor previously or they have not been previously treated for their acute condition. For patients who have received recent previous treatment without relief, CBP rehabilitation care is recommended from the start of treatment [2, 3, 4].

CBP structural rehabilitation is suggested as either three times per week for 12-weeks (36 treatments) or four times per week for 9-weeks (36 treatments), however, the controlled trial data support treatment blocks of 30–40 treatment sessions [7, 8, 9, 10, 15, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55]. An initial patient who has acute or

chronic pains and who has not been treated recently or at all for their current spine issue should be treated for an initial 6–12 sessions to provide pain relief. After signs of relief have occurred, a progress exam should be performed and the patient should be transitioned or ‘graduated’ to CBP corrective care.

CBP treatment occurs in ‘blocks of care.’ Numerous CBP controlled clinical trials (RCTs [[43](#), [44](#), [45](#), [46](#), [47](#), [48](#), [49](#), [50](#), [51](#), [52](#), [53](#), [54](#), [55](#)] and nRCTs [[7](#), [8](#), [9](#), [10](#), [15](#), [42](#)]) provide evidence for spine altering changes to occur in the range of 30–40 treatment sessions; thus, it is the practitioners’ choice to set their protocol within this range (i.e. treatment blocks). The end of each ‘block’ of CBP care requires a progress exam which includes all of the typical assessment procedures as well as a posture and X-ray assessment. Exam results may either dictate the need for further CBP treatment or the recommendation for ‘supportive’ or maintenance care. An initial block of CBP structural rehabilitation will include any acute care provided in the first 2–4 weeks. It is always recommended that ongoing ‘progress exams’ be performed regularly, at

either 4-week or 12 treatment intervals, or as frequently as recommended by each practitioner's regional regulatory board requirements.

CBP does not specifically support 'long-term' care plans. However, based on the data, an adult typically needs 6-months of corrective care (e.g. 72 treatments over 6-months at 3x/week) which is an evidence-based recommendation. Although, any given patient may require a shorted (i.e. 3-month) or longer treatment program based on their initial presenting postural parameters— approximate treatment extrapolations can be made by studying Tables 1–3. There is also support for supportive/maintenance care at a frequency of approximately 2x/month [8, 9, 10].

Study	Journal	Traction method	Traction time	Number of treatments
RCTs				
Moustafa	Sci Reports	Denneroll	20m	30

Moustafa	Heliyon	Denneroll	15–20m	30
Moustafa	J Athl Train	Denneroll	20m	30
Moustafa	APMR	Denneroll	20m	30
Moustafa	EJPRM	Denneroll	20m	30
Moustafa	BFPTCU	Denneroll	20m	36
nRCTs				
Harrison	JMPT	Pope 2-way	20m	38
Harrison	APMR	2way	20m	35
Harrison	JMPT	Ext-comp	10m	60

### **Table 1.**

Summary of cervical lordosis improvement by number of treatments, magnitude correction/treatment and the extrapolation to typical sagittal cervical curve subluxation types and the theoretical treatment number required for their correction to  $-35^{\circ}$  C2-7 ARA.

\*Note: Correction is estimated to achieve  $-35^{\circ}$  of cervical lordosis.

Study	Journal	Traction method	Traction time	Number of treatments
RCTs				
Moustafa	JBMR/JMPT	LET	20m	30
Moustafa	Clin Rehab	LET	20m	30
nRCTs				
Harrison	APMR	LET	20m	36

## Table 2.

Summary of lumbar lordosis improvement by number of treatments, magnitude correction/treatment and the extrapolation to typical sagittal lumbar curve subluxation types and the theoretical treatment number required for their correction to  $-40^\circ$  L1-5 ARA.

\*Note: Correction is estimated to achieve  $-40^\circ$  of lumbar lordosis.

Study	Journal	Traction method	Traction time	Number of treatments
-------	---------	-----------------	---------------	----------------------

nRCTs				
Head trans Harrison	JRRD	Lat trans	20 m	37
Thorax trans Harrison	Eur Sp J	Lat trans	20 m	36

### **Table 3.**

Summary of AP head and thorax lateral translation reduction by number of treatments, magnitude correction/treatment and the extrapolation to larger coronal plane offset subluxations and the theoretical treatment number required for their correction.

Note: Correction is estimated to achieve 0mm of offset.



# 8. Clinical evidence of efficacy

As mentioned, CBP technique has an abundance of clinical evidence supporting its effectiveness in correcting spine deformity and posture [7, 8, 9, 10, 15, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55]. Recently, systematic reviews have summarized the clinical evidence as reported in the published controlled trials on these methods [56, 57]. We summarize the evidence here in four parts: cervical lordosis, lumbar lordosis, lateral translation (pseudo-scoliosis) postures of the head and thorax, and finally, evolving evidence from case reports/series on other important spine deformities including lumbar spondylolisthesis, cervical spondylolisthesis, thoracic hyperkyphosis, thoraco-lumbar junctional kyphosis, thoracic hypokyphosis (straight back syndrome), anterior sagittal balance, lumbar kyphosis (flat back syndrome), lumbar hyperlordosis, post-surgical cervical spine fusion and scoliosis.

## 8.1 Cervical lordosis


A recent systematic review found that of the

RCTs and nRCTs on CBP extension traction methods, a 12–18° improvement in cervical lordosis can be achieved in 10–15 weeks after 30–36 treatment sessions [57]. Most RCTs have used the cervical Denneroll [43, 44, 45, 46, 47, 49, 50], and the three nRCTs all used different CET methods (Table 1) [7, 8, 9].

Table 1 shows the improvement in degrees per treatment as well as theoretical numbers of treatments for various presenting cervical spine subluxations. On average, there appears to be just less than a half degree improvement per treatment session; obviously, there are patients that will have both more correction and less correction than this. Using this estimation as an initial guideline, evidence-based treatment numbers can be predicted. For example, a patient presenting with a cervical kyphosis of 20° would require over 100 treatments to restore the neck to a curve of 35°.


Figures 13 and 14 show the long-term outcomes in patients receiving cervical extension traction versus comparative groups not receiving the traction. The patients restoring lordosis via CBP traction methods

show improved cervical alignment which is maintained at a years' follow-up (Figure 13) whereas, comparative groups receiving various physiotherapeutic treatments less the extension traction do not experience cervical improvement (Figure 13) and also show that any initial pain relief regresses back towards baseline levels after the cessation of treatment (Figure 14). Patient's with improved lordosis retain their initial pain relief a year later (Figure 14). This is alarming as it shows patients receiving various physiotherapeutic treatments who do not improve their cervical lordosis (in hypolordotic patients) will have a future regression of symptoms post-treatment and may be misled by 'apparent treatment efficacy' [5, 57].



## Figure 13.

Data from five RCTs demonstrates patients achieving cervical lordosis improvement (via extension traction) as well as conventional treatments have lordosis improvements that are sustained for 1 year after stopping treatment versus the cervical curve of comparative groups (controls not achieving lordosis improvement) remain unaffected by conventional treatments (weighted averages from five RCTs [44, 45, 47, 49, 50]). \* indicates a significant group difference as specified in each of the five trials; brackets represent weighted standard deviation.



## Figure 14.

Data from five RCTs demonstrates patients achieving cervical lordosis improvement (via extension traction) as well as conventional treatments have pain reductions that are sustained for 1 year after stopping treatment versus

comparative groups (controls not achieving lordosis improvement) who show a regression (increase) of pain intensity towards baseline after stopping treatment (weighted averages from five RCTs [45, 46, 47, 49, 50]). \* indicates a significant group difference as specified in each of the five trials; brackets represent weighted standard deviation.

## 8.2 Lumbar lordosis


A recent systematic review found “Limited but good quality evidence substantiates that the use of extension traction methods in rehabilitation programs definitively increases lumbar hypolordosis” [56]. The authors further stated: “Preliminarily, these studies indicate these methods provide longer-term relief to patients with low back disorders versus conventional rehabilitation approaches tested” [56]. On average, a 7–11° increase in lordosis can be achieved over 10–12 weeks after 30–36 treatment sessions (Table 2).

It must be mentioned that lumbar extension traction is necessary to increase the lumbar lordosis. Importantly, using the data from

published trials [10, 53, 54, 55], one can extrapolate approximate treatment duration (Table 2). As seen, a mild hypolordotic lumbar spine of 30° (L1-L5 ARA) may only require 32–48 treatments, whereas, a flat lumbar curve would require 127–194 treatments to achieve a normal 40° lordosis.

The same trend as observed in patients receiving cervical lordosis correction versus comparative groups not receiving lordosis improvement is seen in the trials on the lumbar spine [5, 56]. Lordosis increase in patients receiving lumbar extension traction is achieved and maintained at 6-months follow-up (Figure 15); these patients also retain their initial pain relief whereas, comparative patient groups not receiving lordosis improvement (Figure 15) lose their initial pain relief by 6-months after cessation of treatment (Figure 16). Again, this is alarming and shows how active low back treatment, although offering transient pain relief, will likely regress after treatment if not receiving concurrent lordosis correction in those suffering from hypolordotic-related LBP [5, 56].


Lumbar lordosis rehabilitation shows  
relative stability at 6-month follow-up  
(n=144 from 2 RCTs)



**Figure 15.**

Data from two RCTs demonstrates patients achieving lumbar lordosis improvement (via extension traction) as well as conventional treatments have lordosis improvements that are sustained for 6-months after stopping treatment versus the lumbar curve of comparative groups (controls not achieving lordosis improvement) remain unaffected by conventional treatments (weighted averages from two RCTs [53, 54]). \* indicates a significant group difference as specified in each of the two trials; brackets represent weighted standard deviation.

Lumbar lordosis rehabilitation leads to better long-term  
pain reduction 6-months post-treatment  
(n= 144 from 2 RCTs)





## Figure 16.

Data from two RCTs demonstrates patients achieving lumbar lordosis improvement (via extension traction) as well as conventional treatments have pain reductions that are sustained for 6-months after stopping treatment versus comparative groups (controls not achieving lordosis improvement) who show a regression (increase) of pain intensity towards baseline after stopping treatment (weighted averages from two RCTs [53, 54]). \* indicates a significant group difference as specified in each of the two trials; brackets represent weighted standard deviation.

## 8.3 AP head and thorax postures

Coronal plane lateral translations of the head and thorax also referred to as ‘pseudo-scoliosis’ each has an nRCT published [15, 42] and many case reports demonstrating its reduction [16, 58, 59, 60, 61, 62, 63]. As discussed earlier, the differentiation from true scoliosis is that the involved vertebrae have minimal to no rotation, whereas, true scoliosis has substantial vertebral rotation (Figure 6). Although the nRCTs demonstrate



(Figure 6). Also, the spinal coupling pattern of a laterally translated body mass (head or thorax) will demonstrate the lower involved spinal region to laterally flex towards the side of the translation and the upper involved spinal region to laterally flex back towards the vertical [35, 36].

Based on the data, a laterally translated body mass can be reduced about 7–8 mm after about 35 treatments. On average, correction of a laterally translated head or thorax can be corrected at about 0.2 mm per treatment, or about 1 mm per five treatments.

Extrapolations of treatment numbers to patient subluxation presentation are shown in Table 3. From the data in each of the nRCTs, an approximate 50% reduction of the initial laterally translated head and thorax postures occurred; therefore, an average patient having an approximate 15 mm translation posture (head or rib cage) requires 6-months of corrective care (approximately 72 treatments). It must also be mentioned that many case reports have demonstrated larger lateral translation postural corrections/reductions with CBP methods in similar time frames [16, 58, 59, 60, 61, 62, 63], thus, these serve as

approximate treatment extrapolations.

## **8.4 Other spine deformities**

It is known that the science for manual therapies is lacking [64]. Therefore, lesser forms of evidence must be considered when evaluating various treatment approaches used to treat various spinal conditions by manual therapists [65, 66]; this includes treatment utilizing CBP methods. We now highlight more recent case studies and series showing structural spinal correction for a variety of relatively common disorders.

### **8.4.1 Lumbar spondylolisthesis**

Fedorchuk et al. [67] reported on an 11 mm reduction (13.3–2.4 mm) of an L4 anterolisthesis in a 69-year old suffering from LBP and leg cramping. Pain relief was achieved after 60 treatments over 45 weeks. This was the first documented report of a reduction of a Grade 2 lumbar spondylolisthesis by CBP methods, as well as any other non-surgical method.

Oakley and Harrison reported on the reduction of multiple retrolistheses from L1-

L4 ranging from 4.5 to 5.9 mm in a 32-year old male with LBP [68]. These were all reduced to within normal (<4.5 mm) after approximately 36 treatments over 14-weeks. A 13-month follow-up indicated the patient remained well and reported no back pain and the corrections had remained stable.

Fedorchuk et al. [69] reported on the reduction of L1 (-6.6 to -1.7 mm) and L2 (-6.1 to -2.0 mm) retrolistheses and an L5 anterolisthesis (+6.8 to -2.5 mm) in a 63-year old female bodybuilder with severe LBP and osteoarthritis. Thirty treatments were given over 10-weeks which resulted in normalizing all spondylolistheses as well as a dramatic reduction in pain and an ability to leg press 60 more pounds in the gym.

Fedorchuk et al. reported the complete reduction of an L3 retrolisthesis and L4 anterolisthesis after 50 treatments over a 7-month period [70]. The patient was 57-years old with severe LBP and sciatica. The L3 retrolisthesis reduced from -5.3 to -1.7 and the L4 anterolisthesis reduced from +5.4 to +1.0 mm. After treatment the patient was able to return to playing hockey and

experienced full resolution of the back pain which had forced him to retire from sport. A 1-year follow-up showed the patient had remained well and maintained the corrections.

### **8.4.2 Cervical spondylolisthesis**

Recently, Fedorchuk et al. present a case series of eight female patients with concomitant cervical hypolordosis, forward head translation and spondylolistheses [71]. All were in motor vehicle collisions, each having at least one, and at most four simultaneous cervical vertebral spondylolistheses ranging in magnitude from >2 mm up to 4.5 mm. All cases experienced a reduction in translational offset of the spondylolistheses, and increase in cervical lordosis and a decrease in forward head translation as well as an increase in spinal canal diameter at the location of the spondylolisthesis after 30 treatment sessions that included cervical extension traction over a duration of 12-weeks. On average, the spondylolistheses reduced by 2.6 mm and there was an average drop in neck disability by 30%.

In another case, Fedorchuk et al. presented a

single case of a 52-year old with chronic neck pain [72]. The patient had a C4 anterolisthesis of 2.4 mm which was reduced to 0.7 mm as well as an increase in cervical lordosis and reduction in forward head translation after 30 treatments over 12-weeks. The patient reported a resolution of their neck pain and stiffness.

### **8.4.3 Thoracic hyperkyphosis**

Thoracic hyperkyphosis is a relatively common subluxation pattern in the aging. Although there is one RCT on CBP methods showing reduction of the deformity, it is yet to be formally published [52]. A systematic review of CBP methods used to reduce thoracic hyperkyphosis was published [73] and summarized the outcomes of several case reports and series [74, 75, 76, 77, 78, 79]. In Table 2 of the Oakley and Harrison review an average 12° reduction in thoracic kyphosis occurred after 32 treatments over 14.5 weeks from a total of 17 patients [52]. The improved posture correlated with reduced pain, disability and improved QOL [52]. Figures 17 and 18 show various CBP mirror image spinal exercises and traction, respectively.



**Figure 17.**  
CBP recommended mirror image exercises for patients with thoracic hyper-kyphosis.



**Figure 18.**



CBP mirror image traction for patients with thoracic hyper-kyphosis.

#### **8.4.4 Thoracolumbar junctional kyphosis**

Thoracolumbar kyphosis is the forward angled spine at the junction of the thoracic and lumbar spine and is associated with chronic LBP (CLBP). Gubbels et al. presented a case of the minimization of pain in a 16-year old female after a 22° reduction of thoracolumbar kyphosis, a 48 mm reduction of posterior sagittal balance, an 11° increase in lumbar lordosis and a 10° increase in sacral inclination [80]. Twenty-four in office treatments were given over an 8-week period with daily home traction resulting in a minimization of back pains.

#### **8.4.5 Thoracic hypokyphosis (straight back syndrome)**

Thoracic spine hypolordosis is termed straight back syndrome (SBS) and is associated with back pains and exertional dyspnea. Fortner et al. [81] reported on an 18-year old male suffering from back pains and exertional dyspnea. Twenty-four treatments over a 9-week period resulted in a

15° increase in thoracic kyphosis, a decrease in pain and improved exertional dyspnea symptoms. A 4-month follow-up showed the patient remained well.

Betz et al. [82] reported the improvement in a 19-year old male who suffered from exertional dyspnea and back pain. Over 12-weeks a 14° increase in thoracic curve was achieved resulting in relief of exertional dyspnea and pain, as well as increases in both the antero-posterior thoracic diameter and the ratio of antero-posterior to transthoracic diameter, both measures critical to the wellbeing of patients with SBS. A 2.75-year follow-up showed the patient remained well.

Fedorchuk et al. [83] reported on a 13° increased thoracic curve in a 26-year old male with back pains and type 1 diabetes. Treatment over 7-weeks included 36 sessions. Back pains reduced and importantly, there was also improvement in blood glucose immediately following the onset of each visit. An improvement in blood glucose averages, percentage of time of blood glucose in a healthy target range, and glycosylated hemoglobin occurred and the patient was



able to reduce their basal insulin need by approximately half after the 7-weeks of care.

Mitchel et al. [84] reported a 10° increase in thoracic curve over 16-weeks in a 33-year old male suffering from exertional dyspnea and back pains. The measured lung capacity improved by 2L, the back pain diminished and the exertional dyspnea resolved. A 7-month follow-up indicated the patient remained well.

#### **8.4.6 Anterior sagittal balance**

Anterior sagittal balance (ASB) is the forward displacement of the upper body over the pelvis. Haas et al. reported on the dramatic 110 mm reduction in ASB in an 87-year old female with CLBP and sciatica [85]. Treatment consisted of 24 in office sessions over an 8-week period. The patient achieved a dramatic reduction of symptoms, improvements in flexibility and orthopedic testing.

Anderson et al. [86] reported on a 91 mm reduction in ASB in a 59-year old male patient suffering from a variety of symptoms associated with Parkinson's disease. Initial treatment involved 28 treatments over

treatment involved 38 treatments over 5 months. The patient experienced significant improvements in multiple postural parameters, gait, balance, hand tremors, low back and knee pains and SF-36 values. A 21-month follow-up showed the patient remained essentially well and most of the initial postural improvements were maintained.

### **8.4.7 Lumbar kyphosis (flat back syndrome)**

Flat back syndrome (FBS) is the anterior translation of the upper body and gross loss (or kyphosis) of the lumbar spine and is associated with high pain and disability. In a case series, Harrison and Oakley describe the significant restoration of lumbar lordosis in two patients suffering from debilitating CLBP from flat back syndrome [87]. One patient had a 50° lordosis improvement in 100 treatments over 20 weeks, the other had a 26° lordosis improvement in 70 treatments over 16.5 weeks. In the discussion section of the report, it was calculated that the treatment costs of the patients receiving CBP treatment versus the projected costs for the surgical procedures recommended to the two patients equated to only 1–8%; the authors

stated “at first 70 or 100 treatments may be criticized as ‘over-treatment,’ however, considering the overall cost-effectiveness and positive patient outcomes, it certainly is not” [87].

#### **8.4.8 Lumbar hyperlordosis**

Although lumbar hypolordosis is the most common lumbar misalignment in those presenting with chronic LBP [10], lumbar hyperlordosis is also seen clinically. CBP methods can be directed at decreasing lumbar lordosis and its typically associated anteriorly rotated pelvis. In a recent case, Oakley et al. [88] presented a case demonstrating the relief of CLBP and hip pains after an 8° reduction in lumbar hyperlordosis, a 5° reduction in pelvic tilt and an accompanying 17 mm reduction of forward sagittal balance. This occurred over a period of 13 months and 73 total treatments.

#### **8.4.9 Post-surgical cervical spine fusion**

Post-surgical cervical spine intervertebral fusion is not a common finding in clinical practice however, it is occasionally

encountered. Many of these patients continue to suffer years after the intervention. Harrison et al. [89] presented a case showing improvement in sagittal postural parameters which corresponded with improved clinical outcome in a 52-year old male. Over a 6-month period, a  $6^\circ$  increase in cervical lordosis was achieved as well as a 13 mm reduction in anterior head translation (AHT). These improvements were maintained at a 2.5-year follow-up.

Fedorchuk et al. [90] also presented a successful outcome in a 43-year old with a C5-6 intersegmental fusion. After 36 treatments over 3-months, there was a  $13^\circ$  increase in cervical lordosis, a 9 mm decrease in AHT and a 5 mm reduction in lateral head translation.

#### **8.4.10 Scoliosis**

Although too large of a topic to address in this chapter, CBP technique has a unique approach in the treatment of scoliosis [3]. CBP methods incorporates the ‘non-commutative property of finite rotation angles under addition’ to ascertain the order of postural movements to be prescribed in the mirror image treatment of this disorder

the minor image treatment of this disorder. Harrison and Oakley described reductions in curve magnitude in five lumbar or thoracolumbar scoliosis patients ranging from 5° to 24° after 18–84 treatments [40]. All patients were female and ranged in age from 19 to 45 years.

Haggard et al. reported a 19° reduction in a thoracolumbar curve in a 15-year old female patient after 24 office treatments over 15-weeks. The patient also performed 45 at home spine blocking sessions as prescribed by the attending chiropractor [41]. The patients LBP and headaches were dramatically improved, and the curve was reduced to 8°.

## 9. Use of X-ray

Use of X-ray for spine analysis is essential for treating spine deformities, including with CBP technique methods. Historically, there has been concerns of carcinogenicity associated with X-ray use. Recently, however, new evidence has come to light showing that anti-X-ray sentiment stemming from the supposed carcinogenicity is based on flawed science [91, 92, 93]. The bottom line is the linear no-threshold (LNT) model used to support radiation risk analysis is not scientific as it is not consistent with current radiobiological data [94, 95, 96, 97, 98].

X-rays and CT scans deliver low-dose radiation doses (<200 mGy), and because of

this they cannot cause cancer. This is because low-dose (versus high-dose) radiation exposures stimulate the adaptive repair systems of the body to repair any damage done [99, 100, 101]. Although this topic is important, it is a much larger issue than the scope of this chapter but many recent reviews have found that X-rays (and CT scans) are not harmful. In fact, after a substantial and critical review of higher quality studies on radiation exposure, Schultz et al. concluded: “The evidence suggests that exposure to multiple CT scans and other sources of low-dose radiation with a cumulative dose up to 100 mSv (approximately 10 scans), and possibly as high as 200 mSv (approximately 20 scans), does not increase cancer risk.” Thus, there should be no hesitation or misunderstanding surrounding X-ray risks. Doctors and patients need to become updated on X-ray safety and not succumb to the traditional carcinogenicity misinformation.

## **10. Conclusion**

CBP technique is a well-studied approach to the structural improvement of spinal disorders. Many spinal disorders with associated pain and functional syndromes have either well characterized or evolving evidence for their treatment by the mirror image approach that underpins CBP methods. The correlation of the spine alignment and postural rotations and translations of posture are of critical importance and unique in the CBP approach.

## **Acknowledgments**

We acknowledge the pioneering work of Dr. Donald D. Harrison.

## **Conflict of interest**

D.E.H. teaches spine rehabilitation methods and sells products related to the treatment of



1  
spine deformities; P.A.O. is a paid consultant  
to CBP.

## Nomenclature

AHT

anterior head translation

ASB

anterior sagittal balance

AP

anterior-to-posterior

ARA

absolute rotation angle

CBP

Chiropractic BioPhysics<sup>®</sup>

CLBP

chronic low back pain

LHT

HPT

Harrison posterior tangent

IVD

intervertebral disc

LBP

low back pain

LNT

linear no-threshold

nRCT

non-randomized controlled trial

QOL

quality of life

PA

posterior-to-anterior

PNF

proprioceptive neuromuscular facilitation

RCT

randomized controlled trial

RRA

relative rotation angle

SEM

standard error of measurement

SBS

straight back syndrome

**INVEST IN YOUR  
HEALTH**

Ads By [tapnative](#)



**1 Teaspoon On An Empty Stomach  
Burns 12 Lbs Of Fat Per Week**

Women Life Wellness





## **A Teaspoon On An Empty Stomach Burns 12 Lbs Of Fat Per Week**

My Health Cycle



## **Randy Jackson: "This Drink Is Like A Powerwash For Your Gut"**

Unify Health Labs



## **1 Teaspoon On An Empty Stomach Can Burn 7Lbs Fat Per Week**



## The 1 Thing You Can Do At Home To Snap Back Sagging Skin

Dermal Medix



## Why Doctors In The Know No Longer Prescribe Metformin

GlucoBurn

# References

1. Harrison DD, Janik TJ, Harrison GR, Troyanovich S, Harrison DE, Harrison SO. Chiropractic biophysics technique: A linear algebra approach to posture in chiropractic. *Journal of Manipulative and Physiological Therapeutics*. 1996;19(8):525-535
2. Harrison DE, Harrison DD, Haas JW. *Structural Rehabilitation of the Cervical Spine*. Evanston, WY: Harrison CBP<sup>®</sup> Seminars, Inc.; 2002
3. Harrison DE, Betz JW, Harrison DD, et al. *CBP Structural Rehabilitation of the Lumbar Spine*. Eagle, ID, USA: Harrison Chiropractic Biophysics Seminars; 2007
4. Oakley PA, Harrison DD, Harrison DE, Haas JW. Evidence-based protocol for structural rehabilitation of the spine and posture: Review of clinical biomechanics of posture (CBP) publications. *Journal of the Canadian Chiropractic Association*. 2005;49(4):270-296
5. Oakley PA, Moustafa IM, Harrison DE.

Restoration of Cervical and Lumbar Lordosis: CBP<sup>®</sup> Methods Overview. In: Bettany-Saltikov J, Kandasamy G, editors. Spinal Deformities in Adolescents, Adults and Older Adults [Internet]. London: IntechOpen; 2019 [cited 2022 Apr 26]. DOI: 10.5772/intechopen.90713

6. Panjabi MM, White AA 3rd, Brand RA Jr. A note on defining body parts configurations. *Journal of Biomechanics*. 1974;7(4):385-387
7. Harrison DD, Jackson BL, Troyanovich S, Robertson G, de George D, Barker WF. The efficacy of cervical extension-compression traction combined with diversified manipulation and drop table adjustments in the rehabilitation of cervical lordosis: A pilot study. *Journal of Manipulative and Physiological Therapeutics*. 1994;17(7):454-464
8. Harrison DE, Cailliet R, Harrison DD, Janik TJ, Holland B. A new 3-point bending traction method for restoring cervical lordosis and cervical manipulation: A nonrandomized clinical controlled trial. *Archives of Physical Medicine and Rehabilitation*



Medicine and Rehabilitation.

2002;83(4):447-453

9. Harrison DE, Harrison DD, Betz JJ, Janik TJ, Holland B, Colloca CJ, et al. Increasing the cervical lordosis with chiropractic biophysics seated combined extension-compression and transverse load cervical traction with cervical manipulation: Nonrandomized clinical control trial. *Journal of Manipulative and Physiological Therapeutics*. 2003;26(3):139-151
10. Harrison DE, Cailliet R, Harrison DD, Janik TJ, Holland B. Changes in sagittal lumbar configuration with a new method of extension traction: Nonrandomized clinical controlled trial. *Archives of Physical Medicine and Rehabilitation*. 2002;83(11):1585-1591
11. CBP NonProfit. [www.cbppnonprofit.com](http://www.cbppnonprofit.com).
12. Beer FP, Johnston ER. *Vector Mechanics for Engineers: Statics and Dynamics*. 4th ed. New York: McGraw-Hill; 1984. p. 95
13. Harrison DD. Abnormal postural permutations calculated as rotations and translations from an ideal normal



upright static spine. In: Sweere J, editor. Chiropractic Family Practice. Gaithersburg, MD: Aspen Publishers; 1992

14. Harrison DE, Betz JW, Cailliet R, Colloca CJ, Harrison DD, Haas JW, et al. Radiographic pseudoscoliosis in healthy male subjects following voluntary lateral translation (side glide) of the thoracic spine. Archives of Physical Medicine and Rehabilitation. 2006;87(1):117-122
15. Harrison DE, Cailliet R, Betz JW, Harrison DD, Colloca CJ, Haas JW, et al. A non-randomized clinical control trial of Harrison mirror image methods for correcting trunk list (lateral translations of the thoracic cage) in patients with chronic low back pain. European Spine Journal. 2005;14(2):155-162
16. Henshaw M, Oakley PA, Harrison DE. Correction of pseudoscoliosis (lateral thoracic translation posture) for the treatment of low back pain: A CBP<sup>®</sup> case report. Journal of Physical Therapy Science. 2018;30(9):1202-1205
17. Harrison DD, Janik TL, Troyanovich SI

17. Harrison DD, Janik TJ, Troyanovich SJ, Holland B. Comparisons of lordotic cervical spine curvatures to a theoretical ideal model of the static sagittal cervical spine. *Spine*. 1996;21(6):667-675
18. Harrison DD, Janik TJ, Troyanovich SJ, Harrison DE, Colloca CJ. Evaluations of the assumptions used to derive an ideal normal cervical spine model. *Journal of Manipulative and Physiological Therapeutics*. 1997;20(4):246-256
19. Harrison DD, Harrison DE, Janik TJ, Cailliet R, Haas JW, Ferrantelli J, et al. Modeling of the sagittal cervical spine as a method to discriminate hypolordosis: Results of elliptical and circular modeling in 72 asymptomatic subjects, 52 acute neck pain subjects, and 70 chronic neck pain subjects. *Spine*. 2004;29:2485-2492
20. Harrison DE, Janik TJ, Harrison DD, Cailliet R, Harmon S. Can the thoracic kyphosis be modeled with a simple geometric shape? The results of circular and elliptical modeling in 80 asymptomatic subjects. *Journal of Spinal Disorders*. 2002;15(3):213-220

21. Harrison DE, Harrison DD, Janik TJ, Cailliet R, Haas JW. Do alterations in vertebral and disc dimensions affect an elliptical model of the thoracic kyphosis? *Spine*. 2003;28(5):463-469
22. Troyanovich SJ, Cailliet R, Janik TJ, Harrison DD, Harrison DE. Radiographic mensuration characteristics of the sagittal lumbar spine from a normal population with a method to synthesize prior studies of lordosis. *Journal of Spinal Disorders*. 1997;10(5):380-386
23. Janik TJ, Harrison DD, Cailliet R, Troyanovich SJ, Harrison DE. Can the sagittal lumbar curvature be closely approximated by an ellipse? *Journal of Orthopaedic Research*. 1998;16(6):766-770
24. Harrison DD, Cailliet R, Janik TJ, Troyanovich SJ, Harrison DE, Holland B. Elliptical modeling of the sagittal lumbar lordosis and segmental rotation angles as a method to discriminate between normal and low back pain subjects. *Journal of Spinal Disorders*

subjects. *Journal of Spinal Disorders*.  
1998;11(5):430-439

25. Harrison DE, Harrison DD, Cailliet R, et al. Cobb method or Harrison posterior tangent method: which to choose for lateral cervical radiographic analysis. *Spine*. 2000;25:2072-2078
26. Harrison DE, Cailliet R, Harrison DD, et al. Reliability of centroid, Cobb, and Harrison posterior tangent methods: Which to choose for analysis of thoracic kyphosis. *Spine*. 2001;26:E227-E234
27. Harrison DE, Harrison DD, Cailliet R, et al. Radiographic analysis of lumbar lordosis: Centroid, Cobb, TRALL, and Harrison posterior tangent methods. *Spine*. 2001;26:E235-E242
28. Harrison DE, Holland B, Harrison DD, et al. Further reliability analysis of the Harrison radiographic line drawing methods: Crossed ICCs for lateral posterior tangents and AP modified-Risser Ferguson. *Journal of Manipulative and Physiological Therapeutics*. 2002;25:93-98

29. McAviney J, Schulz D, Bock R, Harrison DE, Holland B. Determining the relationship between cervical lordosis and neck complaints. *Journal of Manipulative and Physiological Therapeutics*. 2005;28(3):187-193
30. Keller TS, Colloca CJ, Harrison DE, Harrison DD, Janik TJ. Influence of spine morphology on intervertebral disc loads and stresses in asymptomatic adults: Implications for the ideal spine. *The Spine Journal*. 2005;5(3):297-309
31. Harrison DE, Colloca CJ, Harrison DD, Janik TJ, Haas JW, Keller TS. Anterior thoracic posture increases thoracolumbar disc loading. *European Spine Journal*. 2005;14(3):234-242
32. Keller TS, Harrison DE, Colloca CJ, Harrison DD, Janik TJ. Prediction of osteoporotic spinal deformity. *Spine*. 2003;28(5):455-462
33. Harrison DE, Harrison DD, Janik TJ, William Jones E, Cailliet R, Normand M. Comparison of axial and flexural stresses in lordosis and three buckled configurations of the cervical spine.

Configurations of the cervical spine.  
Clinical Biomechanics. 2001;16(4):276-284

34. Harrison DE, Jones EW, Janik TJ, Harrison DD. Evaluation of axial and flexural stresses in the vertebral body cortex and trabecular bone in lordosis and two sagittal cervical translation configurations with an elliptical shell model. Journal of Manipulative and Physiological Therapeutics. 2002;25(6):391-401
35. Harrison DE, Harrison DD, Cailliet R, Janik TJ, Troyanovich SJ. Cervical coupling during lateral head translations creates an S-configuration. Clinical Biomechanics (Bristol, Avon). 2000;15(6):436-440
36. Harrison DE, Cailliet R, Harrison DD, Janik TJ, Troyanovich SJ, Coleman RR. Lumbar coupling during lateral translations of the thoracic cage relative to a fixed pelvis. Clinical Biomechanics (Bristol, Avon). 1999;14(10):704-709
37. Harrison DE, Cailliet R, Harrison DD, Janik TJ. How do anterior/posterior

translations of the thoracic cage affect the sagittal lumbar spine, pelvic tilt, and thoracic kyphosis? *European Spine Journal*. 2002;11(3):287-293

38. Harrison DE, Harrison DD, Haas JW, Oakley PA. *Spinal Biomechanics for Clinicians*. Vol. I. Evanston, WY: Harrison Chiropractic Biophysics Seminars, Inc.; 2003
39. Oakley PA, Cuttler JM, Harrison DE. X-ray imaging is essential for contemporary chiropractic and manual therapy spinal rehabilitation: Radiography increases benefits and reduces risks. *Dose-Response*. 2018;16(2):1559325818781437
40. Harrison DE, Oakley PA. Scoliosis deformity reduction in adults: A CBP<sup>®</sup> Mirror image<sup>®</sup> case series incorporating the 'non-commutative property of finite rotation angles under addition' in five patients with lumbar and thoracolumbar scoliosis. *Journal of Physical Therapy Science*. 2017;29(11):2044-2050
41. Haggard JS, Haggard JB, Oakley PA, Harrison DE. Reduction of progressive

Harrison DE. Reduction of progressive thoracolumbar adolescent idiopathic scoliosis by chiropractic biophysics<sup>®</sup> (CBP<sup>®</sup>) mirror image<sup>®</sup> methods following failed traditional chiropractic treatment: A case report. *Journal of Physical Therapy Science*. 2017;29(11):2062-2067

42. Harrison DE, Cailliet R, Betz J, Haas JW, Harrison DD, Janik TJ, et al. Conservative methods for reducing lateral translation postures of the head: A nonrandomized clinical control trial. *Journal of Rehabilitation Research and Development*. 2004;41(4):631-639
43. Moustafa IM, Diab AA, Hegazy F, Harrison DE. Demonstration of central conduction time and neuroplastic changes after cervical lordosis rehabilitation in asymptomatic subjects: A randomized, placebo-controlled trial. *Scientific Reports*. 2021;11(1):15379
44. Moustafa IM, Diab A, Shousha T, Harrison DE. Does restoration of sagittal cervical alignment improve cervicogenic headache pain and disability: A 2-year pilot randomized



disability. A 2-year pilot randomized controlled trial. *Heliyon*. 2021;7(3):e06467

45. Moustafa I, Youssef ASA, Ahbouch A, Harrison DE. Demonstration of autonomic nervous function and cervical sensorimotor control after cervical lordosis rehabilitation: A randomized controlled trial. *Journal of Athletic Training*. 2021;56(3):10
46. Moustafa IM, Diab AA, Hegazy F, Harrison DE. Does improvement towards a normal cervical sagittal configuration aid in the management of cervical myofascial pain syndrome: A 1-year randomized controlled trial. *BMC Musculoskeletal Disorders*. 2018;19(1):396
47. Moustafa IM, Diab AA, Harrison DE. The effect of normalizing the sagittal cervical configuration on dizziness, neck pain, and cervicocephalic kinesthetic sensibility: A 1-year randomized controlled study. *European Journal of Physical and Rehabilitation Medicine*. 2017;53(1):57-71

48. Moustafa IM, Diab AAM, Hegazy FA, Harrison DE. Does rehabilitation of cervical lordosis influence sagittal cervical spine flexion extension kinematics in cervical spondylotic radiculopathy subjects? *Journal of Back and Musculoskeletal Rehabilitation*. 2017;30(4):937-941
49. Moustafa IM, Diab AA, Taha S, Harrison DE. Addition of a sagittal cervical posture corrective orthotic device to a multimodal rehabilitation program improves short- and long-term outcomes in patients with discogenic cervical radiculopathy. *Archives of Physical Medicine and Rehabilitation*. 2016;97(12):2034-2044
50. Moustafa IM. Does improvement towards a normal cervical configuration aid in the management of fibromyalgia. A randomized controlled trial. *Bulletin of Faculty of Pharmacy, Cairo University*. 2013;18(2):29-41
51. Moustafa IM, Diab AM, Ahmed A, Harrison DE. The efficacy of cervical lordosis rehabilitation for nerve root function, pain, and segmental motion in

function, pain, and segmental motion in cervical spondylotic radiculopathy. *Physiotherapy*. 2011;97(supplement):846-847

52. Moustafa IM, Walton LM, Raigangir V, Shousha TM, Harrison D. Reduction of posture hyperkyphosis improves short- and long-term outcomes in patients with neck pain. Abstract: *International Journal of Orthopaedic & Sports Physical Therapy*. 2020;50(1):CSM143
53. Diab AAM, Moustafa IM. The efficacy of lumbar extension traction for sagittal alignment in mechanical low back pain: A randomized trial. *Journal of Back and Musculoskeletal Rehabilitation*. 2013;26(2):213-220
54. Moustafa IM, Diab AA. Extension traction treatment for patients with discogenic lumbosacral radiculopathy: A randomized controlled trial. *Clinical Rehabilitation*. 2012;27(1):51-62
55. Diab AA, Moustafa IM. Lumbar lordosis rehabilitation for pain and lumbar segmental motion in chronic mechanical low back pain: A randomized trial.

Journal of Manipulative and  
Physiological Therapeutics.  
2012;35(4):246-253

56. Oakley PA, Ehsani NN, Moustafa IM, Harrison DE. Restoring lumbar lordosis: A systematic review of controlled trials utilizing Chiropractic Bio Physics<sup>®</sup> (CBP<sup>®</sup>) non-surgical approach to increasing lumbar lordosis in the treatment of low back disorders. Journal of Physical Therapy Science. 2020;32(9):601-610
57. Oakley PA, Ehsani NN, Moustafa IM, Harrison DE. Restoring cervical lordosis by cervical extension traction methods in the treatment of cervical spine disorders: A systematic review of controlled trials. Journal of Physical Therapy Science. 2021;33(10):784-794.
58. Haas JW, Oakley PA, Harrison DE. Cervical pseudo-scoliosis reduction and alleviation of dystonia symptoms using chiropractic BioPhysics<sup>®</sup> (CBP<sup>®</sup>) technique: A case report with a 1.5-year follow-up. The Journal of Contemporary Chiropractic. 2019;2:131-137

59. Jaeger JO, Oakley PA, Moore RR, Ruggeroli EP, Harrison DE. Resolution of temporomandibular joint dysfunction (TMJD) by correcting a lateral head translation posture following previous failed traditional chiropractic therapy: A CBP<sup>®</sup> case report. *Journal of Physical Therapy Science*. 2018;30(1):103-107
60. Oakley PA, Harrison DE. Alleviation of pain and disability in a post-surgical C4-C7 total fusion patient after reducing a lateral head translation (side shift) posture: A CBP<sup>®</sup> case report with a 14 year follow-up. *Journal of Physical Therapy Science*. 2018;30(7):952-957
61. Berry RH, Oakley P, Harrison D. Alleviation of radiculopathy by structural rehabilitation of the cervical spine by correcting a lateral head translation posture (-TxH) using Berry translation traction as a part of CBP methods: A case report. *The Chiropractic Journal of Australia*. 2017;45(1):63-72
62. Berry RH, Oakley PA, Harrison DE.

Alleviation of chronic headaches by correcting lateral head translation posture (-TxH) using Chiropractic Biophysics & Berry Translation Traction. *Annals of Vertebral Subluxation Research*. 2017;(1-2):87-92.

63. Oakley PA, Berry RH, Harrison DE. A structural approach to the postsurgical laminectomy case. *Journal of Vertebral Subluxation Research*. 2007;(March 19):1-7.
64. Nuckols TK, Lim YW, Wynn BO, et al. Rigorous development does not ensure that guidelines are acceptable to a panel of knowledgeable providers. *Journal of General Internal Medicine*. 2008;23(1):37-44
65. Rome P, Waterhouse JD. An evidence-based narrative of the evidence-base concept. *Asia-Pacific Chiropractic Journal*. 2020;1:004.  
<https://doi.org/10.46323/2021004>
66. Ebrall P, Doyle M. The value of case reports as clinical evidence. *The Chiropractic Journal of Australia*. 2020;47(1):20-42

67. Fedorchuk C, Lightstone DF, McRae C, Kaczor D. Correction of grade 2 spondylolisthesis following a non-surgical structural spinal rehabilitation protocol using lumbar traction: A case study and selective review of literature. *Journal of Radiology Case Reports*. 2017;11(5):13-26
68. Oakley PA, Harrison DE. Correction of multilevel lumbar retrolistheses by non-surgical extension traction procedures in a patient with congenital fusion of L5-S1: A CBP<sup>®</sup> case report with a 13-month follow-up. *The Journal of Contemporary Chiropractic*. 2020;3(1):137-142
69. Fedorchuk C, Haugen H. Reduction in three levels of lumbar degenerative spondylolisthesis following chiropractic care: A case report & review of the literature. *Annals of Vertebral Subluxation Research*. 2020;(Dec. 3):165-170.
70. Fedorchuk CA, Lightstone DF, Oakley PA, Harrison DE. Correction of a double spondylolisthesis of the lumbar spine

utilizing chiropractic biophysics<sup>®</sup>  
technique: A case report with 1-year  
follow-up. Journal of Physical Therapy  
Science. 2021;33(1):89-93

71. Fedorchuk C, Lightstone DF, DeVon CR, Katz E, Wilcox J. Improvements in cervical spinal canal diameter and neck disability following correction of cervical lordosis and cervical spondylolistheses using chiropractic BioPhysics technique: A case series. Journal of Radiology Case Reports. 2020;14(4):21-37
72. Fedorchuk C, Lightstone D. Reduction in cervical anterolisthesis & pain in a 52-year-old female using chiropractic biophysics<sup>®</sup> technique: A case study and selective review of literature. Annals of Vertebral Subluxation Research. 2016;3:118-124
73. Oakley PA, Harrison DE. Reducing thoracic Hyperkyphosis subluxation deformity: A systematic review of chiropractic BioPhysics<sup>®</sup> methods employed in its structural improvement. The Journal of Contemporary



Chiropractic. 2018;1(1):59-66

74. Fortner MO, Oakley PA, Harrison DE. Alleviation of chronic spine pain and headaches by reducing forward head posture and thoracic hyperkyphosis: A CBP<sup>®</sup> case report. Journal of Physical Therapy Science. 2018;30(8):1117-1123
75. Oakley PA, Jaeger JO, Brown JE, Polatis TA, Clarke JG, Whittler CD, et al. The CBP<sup>®</sup> mirror image<sup>®</sup> approach to reducing thoracic hyperkyphosis: A retrospective case series of 10 patients. Journal of Physical Therapy Science. 2018;30(8):1039-1045
76. Fortner MO, Oakley PA, Harrison DE. Treating 'slouchy' (hyperkyphosis) posture with chiropractic biophysics<sup>®</sup>: A case report utilizing a multimodal mirror image<sup>®</sup> rehabilitation program. Journal of Physical Therapy Science. 2017;29(8):1475-1480
77. Miller JE, Oakley PA, Levin SB, Harrison DE. Reversing thoracic hyperkyphosis: A case report featuring mirror image<sup>®</sup> thoracic extension rehabilitation

thoracic extension rehabilitation.  
Journal of Physical Therapy Science.  
2017;29(7):1264-1267

78. Fedorchuk C, Snow E. Reduction in thoracic hyperkyphosis with increased peak expiratory flow (PEF), forced expiratory volume (FEV) and SF-36 scores following CBP protocols in asymptomatic patients: A case series. Annals of Vertebral Subluxation Research. 2017;(Oct. 12):189-200.
79. Jaeger JO, Oakley PA, Colloca CJ, et al. Non-surgical reduction of thoracic hyper-kyphosis in a 24-year old music teacher utilizing chiropractic biophysics<sup>®</sup> technique. British Journal of Medicine and Medical Research. 2016;11:1-9
80. Gubbels CM, Werner JT, Oakley PA, Harrison DE. Reduction of thoracolumbar junctional kyphosis, posterior sagittal balance, and increase of lumbar lordosis and sacral inclination by chiropractic BioPhysics<sup>®</sup> methods in an adolescent with back pain: A case report. Journal of Physical Therapy Science. 2019;21(10):820-842

81. Fortner MO, Oakley PA, Harrison DE. Chiropractic biophysics management of straight back syndrome and exertional dyspnea: A case report with follow-up. *The Journal of Contemporary Chiropractic*. 2019;2:115-122
82. Betz JW, Oakley PA, Harrison DE. Relief of exertional dyspnea and spinal pains by increasing the thoracic kyphosis in straight back syndrome (thoracic hypokyphosis) using CBP<sup>®</sup> methods: A case report with long-term follow-up. *Journal of Physical Therapy Science*. 2018;30(1):185-189
83. Fedorchuk C, Lightstone DF, Comer RD, Weiner MT, McCoy M. Improved glycosylated hemoglobin, hyperglycemia, and quality of life following thoracic hypokyphosis vertebral subluxation correction using Chiropractic BioPhysics<sup>®</sup>: A prospective case report. *Journal of Diabetes & Metabolism*. 2018;9(10):1-10
84. Mitchell JR, Oakley PA, Harrison DE. Nonsurgical correction of straight back

syndrome (thoracic hypokyphosis), increased lung capacity and resolution of exertional dyspnea by thoracic hyperkyphosis mirror image<sup>®</sup> traction: A CBP<sup>®</sup> case report. *Journal of Physical Therapy Science*. 2017;29(11):2058-2061

85. Haas JW, Harrison DE, Oakley PA. Non-surgical reduction in anterior sagittal balance subluxation and improvement in overall posture in a geriatric suffering from low back pain and sciatica: A CBP<sup>®</sup> case report. *The Journal of Contemporary Chiropractic*. 2020;3(1):45-50
86. Anderson JM, Oakley PA, Harrison DE. Improving posture to reduce the symptoms of Parkinson's: A CBP<sup>®</sup> case report with a 21 month follow-up. *Journal of Physical Therapy Science*. 2019;31(2):153-158
87. Harrison DE, Oakley PA. Non-operative correction of flat back syndrome using lumbar extension traction: A CBP<sup>®</sup> case series of two. *Journal of Physical Therapy Science*. 2018;30(8):1131-1137

88. Oakley PA, Ehsani NN, Harrison DE. Non-surgical reduction of lumbar hyperlordosis, forward sagittal balance and sacral tilt to relieve low back pain by Chiropractic BioPhysics® methods: A case report. *Journal of Physical Therapy Science*. 2019;31(10):860-864
89. Harrison DE, Oakley PA, Betz JW. Anterior head translation following cervical fusion-a probable cause of post-surgical pain and impairment: A CBP® case report. *Journal of Physical Therapy Science*. 2018;30(2):271-276
90. Fedorchuk C, Lightstone DF, Andino H. Failed neck surgery: Improvement in neck pain, migraines, energy levels, and performance of activities of daily living following subluxation correction using Chiropractic Biophysics® Technique: A case study. *Annals of Vertebral Subluxation Research*. 2017;(May 18):93-100
91. Calabrese EJ. Cancer risk assessment foundation unraveling: New historical evidence reveals that the US National

Academy of Sciences (US NAS),  
biological effects of atomic radiation  
(BEAR) committee genetics panel  
falsified the research record to promote  
acceptance of the LNT. Archives of  
Toxicology. 2015;89(4):649-650

92. Calabrese EJ. An abuse of risk  
assessment: How regulatory agencies  
improperly adopted LNT for cancer risk  
assessment. Archives of Toxicology.  
2015;89(4):647-648
93. Calabrese EJ. On the origins of the linear  
no-threshold (LNT) dogma by means of  
untruths, artful dodges and blind faith.  
Environmental Research. 2015;142:432-  
442
94. Scott BR, Sanders CL, Mitchel REJ,  
Boreham DR. CT scans may reduce  
rather than increase risk of cancer. The  
Journal of the American Physicians and  
Surgeons. 2008;13(1):8-11
95. Lemon JA, Phan N, Boreham DR. Single  
CT scan prolongs survival by extending  
cancer latency in Trp53 heterozygous  
mice. Radiation Research.  
2017;188(4.2):505-511

96. Lemon JA, Phan N, Boreham DR. Multiple CT scans extend lifespan by delaying cancer progression in cancer-prone mice. *Radiation Research*. 2017;188(4.2):495-504
97. Cuttler JM. Application of low doses of ionizing radiation in medical therapies. Dose-Response. 2020;18(1):1559325819895739
98. Calabrese EJ, Dhawan G, Kapoor R, Kozumbo WJ. Radiotherapy treatment of human inflammatory diseases and conditions: Optimal dose. *Human & Experimental Toxicology*. 2019;38(8):888-898
99. Pollycove M, Feinendegen LE. Radiation-induced versus endogenous DNA damage: Possible effect of inducible protective responses in mitigating endogenous damage. *Human and Experimental Toxicology*. 2003;22(6):290-306
100. Pollycove M. Radiobiological basis of low-dose irradiation in prevention and therapy of cancer. *Dose-Response*

therapy of cancer: Dose response.  
2006;5(1):26-38

101. Feinendegen LE, Cuttler JM. Biological effects from low doses and dose rates of ionizing radiation: Science in the service of protecting humans, a synopsis. Health Physics. 2018;114(6):623-626

---

## WRITTEN BY

**Deed E. Harrison and Paul A. Oakley**

Submitted: October 13th, 2021 , Reviewed: January 16th, 2022 , Published: May 19th, 2022



**REGISTER TO DOWNLOAD FOR  
FREE**



**Share**



**Cite**



properly cited.

CONTINUE READING FROM THE SAME BOOK

[View All](#)



## CHAPTER 9

### Complementary Treatment for Women with Breast Canc...

By Guida Veiga and Graça Duarte Santos

105 DOWNLOADS



## Complementary Therapies

Edited by Mario Bernardo-Filho

PUBLISHED: JULY 6TH, 2022



[Home](#)

[Careers](#)

[News](#)

[About](#)

[Contact](#)

[Publish](#)



Headquarters

Phone: +44 20 8089 5702

IntechOpen Limited

5 Princes Gate Court,

London, SW7 2QJ,

UNITED KINGDOM

Search



[AUTHOR PANEL SIGN IN](#)

---

[Terms and Conditions](#)

[Privacy Policy](#)

[Customer Complaints](#)

© 2022 IntechOpen. All rights reserved.